arXiv:0709.0959v3 [math.AT] 10 Aug 2008

THE MORSE-BOTT INEQUALITIES VIA DYNAMICAL SYSTEMS

AUGUSTIN BANYAGA AND DAVID E. HURTUBISE

Abstract. Let $f : M \to \mathbb{R}$ be a Morse-Bott function on a compact smooth finite dimensional manifold M. The polynomial Morse inequalities and an explicit perturbation of f defined using Morse functions f_j on the critical submanifolds C_j of f show immediately that $MB_t(f) = P_t(M) + (1+t)R(t)$, where $MB_t(f)$ is the Morse-Bott polynomial of f and $P_t(M)$ is the Poincaré polynomial of M. We prove that $R(t)$ is a polynomial with nonnegative integer coefficients by showing that the number of gradient flow lines of the perturbation of f between two critical points $p, q \in C_j$ of relative index one coincides with the number of gradient flow lines between p and q of the Morse function f_j. This leads to a relationship between the kernels of the Morse-Smale-Witten boundary operators associated to the Morse functions f_j and the perturbation of f. This method works when M and all the critical submanifolds are oriented or when $\mathbb{Z}_2$ coefficients are used.

1. Introduction

Let $h : M \to \mathbb{R}$ be a Morse function on a compact smooth manifold of dimension m. The Morse inequalities say that

$$\nu_n - \nu_{n-1} + \cdots + (-1)^n \nu_0 \geq b_n - b_{n-1} + \cdots + (-1)^n b_0$$

for all $n = 0, \ldots, m$ (with $\geq$ an equality when $n = m$), where ν_k is the number of critical points of h of index k and b_k is the k^{th} Betti number of M for all k. These inequalities follow from the fact that the Morse function h determines a CW-complex X whose cellular homology is isomorphic to the singular homology of M (the k-cells of X are in bijective correspondence with the critical points of index k).

In [1414, 14] Witten introduced the idea that the Morse inequalities can be studied by deforming the de Rham differential on differential forms by the differential of the Morse function. This led him to consider a chain complex whose chains are generated by the critical points of the Morse function and whose differential is defined by counting the gradient flow lines between critical points of relative index one (the so-called Morse-Smale-Witten chain complex). See also [99, 9]. The homology of this complex is called the "Morse homology", and the Morse Homology Theorem asserts that the Morse homology is isomorphic to the singular homology (see [33, 3] and [1313, 13]). For an excellent exposition of Witten's ideas see [1111, 11].

2000 *Mathematics Subject Classification.* Primary: 57R70 Secondary: 37D15.

On one hand Bismut, and on the other hand, Helffer and Sjöstrand have given rigorous mathematical derivations of Witten's analytical ideas [66, 6][1010, 10]. In this way, they proved the Morse inequalities and the Morse-Bott inequalities, which relate the Betti numbers of M and the Betti numbers of the critical submanifolds of a Morse-Bott function on M, without using the Morse Homology Theorem.

In this paper, we present a proof of the Morse-Bott inequalities using ideas from dynamical systems. The proof uses the Morse inequalities, the Morse Homology Theorem, and an explicit perturbation technique that produces a Morse-Smale function h arbitrarily close to a given Morse-Bott function f [11, 1] [22, 2]. Roughly, the perturbation h of f is constructed as follows. We first fix a Riemannian metric on M and apply the Kupka-Smale Theorem to choose Morse-Smale functions f_j on the critical submanifolds C_j for $j = 1, \ldots, l$. Next, we extend the Morse-Smale functions f_j to tubular neighborhoods T_j of the critical submanifolds, and we define

$$h = f + \varepsilon \left(\sum_{j=1}^{l} \rho_j f_j \right)$$

where ρ_j is a bump function on T_j and $\varepsilon > 0$. We then apply a well-known folk theorem (whose proof can be found in Section 2.12 of [11, 1]) which says that we can perturb the Riemannian metric on M outside of the union of the tubular neighborhoods T_j for $j = 1, \ldots, l$ so that h satisfies the Morse-Smale transversality condition with respect to the perturbed metric.

Once we know that h and f_j for $j = 1, \ldots, l$ satisfy the Morse-Smale transversality condition, we can compare the Morse-Smale-Witten chain complex of h to the Morse-Smale-Witten chain complexes of f_j for $j = 1, \ldots, l$. This allows us to show that the coefficients of the polynomial $R(t)$ in Theorem 8 are nonnegative. The proof works when the manifold M and all the critical submanifolds are oriented or when $\mathbb{Z}_2$ coefficients are used.

]cite.BotNon777]cite.BotLec888]cite.BanMor555]cite.JiaMor121212

2. The Morse-Smale-Witten chain complex

In this section we briefly recall the construction of the Morse-Smale-Witten chain complex and the Morse Homology Theorem. For more details see [33, 3].

Let $Cr(f) = \{p \in M |\, df_p = 0\}$ denote the set of critical points of a smooth function $f : M \to \mathbb{R}$ on a smooth m-dimensional manifold M. A critical point $p \in Cr(f)$ is said to be **nondegenerate** if and only if the Hessian $H_p(f)$ is nondegenerate. The **index** λ_p of a nondegenerate critical point p is the dimension of the subspace of T_pM where $H_p(f)$ is negative definite. If all the critical points of f are non-degenerate, then f is called a **Morse function**.

If $f : M \to \mathbb{R}$ is a Morse function on a finite dimensional compact smooth Riemannian manifold (M, g), then the **stable manifold** $W^s(p)$ and the **unstable manifold**

$W^u(p)$ of a critical point $p \in Cr(f)$ are defined to be

$$\begin{aligned} W^s(p) &= \{x \in M | \lim_{t\to\infty} \varphi_t(x) = p\} \\ W^u(p) &= \{x \in M | \lim_{t\to-\infty} \varphi_t(x) = p\} \end{aligned}$$

where φ_t is the 1-parameter group of diffeomorphisms generated by minus the gradient vector field, i.e. $-\nabla f$. The index of p coincides with the dimension of $W^u(p)$. The Stable/Unstable Manifold Theorem for a Morse Function says that the tangent space at p splits as

$$T_pM = T_p^sM \oplus T_p^uM$$

where the Hessian is positive definite on $T_p^sM \overset{def}{=} T_pW^s(p)$ and negative definite on $T_p^uM \overset{def}{=} T_pW^u(p)$. Moreover, the stable and unstable manifolds of p are surjective images of smooth embeddings

$$\begin{aligned} E^s : T_p^sM &\to W^s(p) \subseteq M \\ E^u : T_p^uM &\to W^u(p) \subseteq M. \end{aligned}$$

Hence, $W^s(p)$ is a smoothly embedded open disk of dimension $m - \lambda_p$, and $W^u(p)$ is a smoothly embedded open disk of dimension λ_p.

If the stable and unstable manifolds of a Morse function $f : M \to \mathbb{R}$ all intersect transversally, then the function f is called **Morse-Smale**. Note that for a given Morse function $f : M \to \mathbb{R}$ one can choose a Riemannian metric on M so that f is Morse-Smale with respect to the chosen metric (see Theorem 2.20 of [11, 1]). Moreover, if f is Morse-Smale then $W(q,p) = W^u(q) \cap W^s(p)$ is an embedded submanifold of M of dimension $\lambda_q - \lambda_p$, and when $\lambda_q = \lambda_p + 1$ one can use Palis' λ-Lemma to prove that the number of gradient flow lines from q to p is finite.

If we assume that M is oriented and we choose an orientation for each of the unstable manifolds of f, then there is an induced orientation on the stable manifolds. Thus, we can define an integer $n(q,p)$ associated to any two critical points p and q of relative index one by counting the number of gradient flow lines from q to p with signs determined by the orientations. If M is not orientable, then we can still define $n(q,p) \in \mathbb{Z}_2$ by counting the number of gradient flow lines from q to p mod 2.

The **Morse-Smale-Witten chain complex** is defined to be the chain complex $(C_*(f), \partial_*)$ where $C_k(f)$ is the free abelian group generated by the critical points q of index k (tensored with $\mathbb{Z}_2$ when $n(q,p)$ is defined as an element of $\mathbb{Z}_2$) and the boundary operator $\partial_k : C_k(f) \to C_{k-1}(f)$ is given by

$$\partial_k(q) = \sum_{p \in Cr_{k-1}(f)} n(q,p)p.$$

Theorem 1 (Morse Homology Theorem)**.** *The pair $(C_*(f), \partial_*)$ is a chain complex. If M is orientable and the boundary operator is defined with $n(q,p) \in \mathbb{Z}$, then the homology of $(C_*(f), \partial_*)$ is isomorphic to the singular homology $H_*(M;\mathbb{Z})$. If the* *boundary operator is defined with* $n(q,p) \in \mathbb{Z}_2$*, then the homology of* $(C_*(f), \partial_*)$ *is isomorphic to the singular homology* $H_*(M;\mathbb{Z}_2)$*.*

Note that the Morse Homology Theorem implies that the homology of $(C_*(f), \partial_*)$ is independent of the Morse-Smale function $f : M \to \mathbb{R}$, the Riemannian metric, and the orientations.

3. The Morse Inequalities

Let M be a compact smooth manifold of dimension m. When M is orientable we define the k^{th} **Betti number** of M, denoted b_k, to be the rank of the k^{th} homology group $H_k(M;\mathbb{Z})$ modulo its torsion subgroup. When M is not orientable we define $b_k = \dim H_k(M;\mathbb{Z}_2)$. Let $f : M \to \mathbb{R}$ be a Morse function on M, and let ν_k be the number of critical points of f of index k for all $k = 0, \ldots, m$. As a consequence of the Morse Homology Theorem we have

$$\nu_k \geq b_k$$

for all $k = 0, \ldots, m$ since $\nu_k = \text{rank } C_k(f)$ and $H_k(M;\mathbb{Z})$ (or $H_k(M;\mathbb{Z}_2)$) is a quotient of $C_k(f)$. These inequalities are known as the **weak Morse inequalities**.

Definition 2. *The* ***Poincaré polynomial*** *of* M *is defined to be*

$$P_t(M) = \sum_{k=0}^{m} b_k t^k,$$

and the ***Morse polynomial*** *of* f *is defined to be*

$$M_t(f) = \sum_{k=0}^{m} \nu_k t^k.$$

The Morse inequalities stated in the introduction are known as the **strong Morse inequalities**. The strong Morse inequalities are equivalent to the following **polynomial Morse inequalities**. (For a proof see Lemma 3.43 of [33,3].) It is this version of the Morse inequalities that we generalize to Morse-Bott functions.

Theorem 3 (Polynomial Morse Inequalities)**.** *For any Morse function* $f : M \to \mathbb{R}$ *on a smooth manifold* M *we have*

$$M_t(f) = P_t(M) + (1+t)R(t)$$

where $R(t)$ *is a polynomial with non-negative integer coefficients. That is,* $R(t) = \sum_{k=0}^{m-1} r_k t^k$ *where* $r_k \in \mathbb{Z}$ *satisfies* $r_k \geq 0$ *for all* $k = 0, \ldots, m-1$*.*

Although Theorem 3 and its proof are standard facts, we give here a detailed proof using the Morse-Smale-Witten chain complex because this proof gives an explicit formula $r_k = \nu_{k+1} - z_{k+1}$ for the coefficients of the polynomial $R(t)$ which we will use in the proof of Theorem 8.

Proof of Theorem 3: Let $f : M \to \mathbb{R}$ be a Morse function on a finite dimensional compact smooth manifold M, and choose a Riemannian metric on M for which f is a Morse-Smale function. Let $C_k(f)$ denote the k^{th} chain group in the Morse-Smale-Witten chain complex of f (with coefficients in either $\mathbb{Z}$ or $\mathbb{Z}_2$), and let $\partial_k : C_k(f) \to C_{k-1}(f)$ denote the k^{th} Morse-Smale-Witten boundary operator. The rank of $C_k(f)$ is equal to the number ν_k of critical points of f of index k, and by the Morse Homology Theorem (Theorem 1) the rank of $H_k(C_*(f), \partial_*)$ is equal to b_k, the k^{th} Betti number of M for all $k = 0, \ldots, m$.

Let $z_k = \text{rank ker } \partial_k$ for all $k = 0, \ldots, m$. The exact sequence

$$0 \to \ker \partial_k \to C_k(f) \xrightarrow{\partial_k} \text{im } \partial_k \to 0$$

implies that $\nu_k = z_k + \text{rank im } \partial_k$ for all $k = 0, \ldots, m$, and

$$0 \to \text{im } \partial_{k+1} \to \ker \partial_k \to H_k(C_*(f), \partial_*) \to 0$$

implies that $b_k = z_k - \text{rank im } \partial_{k+1}$ for all $k = 0, \ldots, m$. Hence,

$$\begin{aligned}
M_t(f) - P_t(M) &= \sum_{k=0}^{m} \nu_k t^k - \sum_{k=0}^{m} b_k t^k \\
&= \sum_{k=0}^{m} (z_k + \text{rank im } \partial_k) t^k - \sum_{k=0}^{m} (z_k - \text{rank im } \partial_{k+1}) t^k \\
&= \sum_{k=0}^{m} (\text{rank im } \partial_k + \text{rank im } \partial_{k+1}) t^k \\
&= \sum_{k=0}^{m} (\nu_k - z_k + \nu_{k+1} - z_{k+1}) t^k \\
&= \sum_{k=0}^{m} (\nu_k - z_k) t^k + \sum_{k=0}^{m-1} (\nu_{k+1} - z_{k+1}) t^k \\
&= t \sum_{k=1}^{m} (\nu_k - z_k) t^{k-1} + \sum_{k=1}^{m} (\nu_k - z_k) t^{k-1} \quad (\text{since } \nu_0 = z_0) \\
&= (t+1) \sum_{k=1}^{m} (\nu_k - z_k) t^{k-1}.
\end{aligned}$$

Therefore, $M_t(f) = P_t(M) + (1+t)R(t)$ where $R(t) = \sum_{k=0}^{m-1} (\nu_{k+1} - z_{k+1}) t^k$. Note that $\nu_{k+1} - z_{k+1} \geq 0$ for all $k = 0, \ldots, m-1$ because z_{k+1} is the rank of a subgroup of $C_{k+1}(f)$ and $\nu_{k+1} = \text{rank } C_{k+1}(f)$.

$\square$

4. The Morse-Bott Inequalities

Let $f : M \to \mathbb{R}$ be a smooth function whose critical set $\mathrm{Cr}(f)$ contains a submanifold C of positive dimension. Pick a Riemannian metric on M and use it to split $T_*M|_C$ as

$$T_*M|_C = T_*C \oplus \nu_*C$$

where T_*C is the tangent space of C and ν_*C is the normal bundle of C. Let $p \in C$, $V \in T_pC$, $W \in T_pM$, and let $H_p(f)$ be the Hessian of f at p. We have

$$H_p(f)(V, W) = V_p \cdot (\tilde{W} \cdot f) = 0$$

since $V_p \in T_pC$ and any extension of W to a vector field $\tilde{W}$ satisfies $df(\tilde{W})|_C = 0$. Therefore, the Hessian $H_p(f)$ induces a symmetric bilinear form $H_p^\nu(f)$ on ν_pC.

Definition 4. *A smooth function $f : M \to \mathbb{R}$ on a smooth manifold M is called a* ***Morse-Bott function*** *if and only if the set of critical points $\mathrm{Cr}(f)$ is a disjoint union of connected submanifolds and for each connected submanifold $C \subseteq \mathrm{Cr}(f)$ the bilinear form $H_p^\nu(f)$ is non-degenerate for all $p \in C$.*

Often one says that the Hessian of a Morse-Bott function f is non-degenerate in the direction normal to the critical submanifolds.

For a proof of the following lemma see Section 3.5 of [33, 3] or [44, 4].

Lemma 5 (Morse-Bott Lemma). *Let $f : M \to \mathbb{R}$ be a Morse-Bott function and $C \subseteq \mathrm{Cr}(f)$ a connected component. For any $p \in C$ there is a local chart of M around p and a local splitting $\nu_*C = \nu_*^-C \oplus \nu_*^+C$, identifying a point $x \in M$ in its domain to (u, v, w) where $u \in C$, $v \in \nu_*^-C$, $w \in \nu_*^+C$, such that within this chart f assumes the form*

$$f(x) = f(u, v, w) = f(C) - |v|^2 + |w|^2.$$

Definition 6. *Let $f : M \to \mathbb{R}$ be a Morse-Bott function on a finite dimensional smooth manifold M, and let C be a critical submanifold of f. For any $p \in C$ let λ_p denote the index of $H_p^\nu(f)$. This integer is the dimension of ν_p^-C and is locally constant by the preceding lemma. If C is connected, then λ_p is constant throughout C and we call $\lambda_p = \lambda_C$ the* ***Morse-Bott index*** *of the connected critical submanifold C.*

Let $f : M \to \mathbb{R}$ be a Morse-Bott function on a finite dimensional compact smooth manifold, and assume that

$$\mathrm{Cr}(f) = \coprod_{j=1}^{l} C_j,$$

where $C_1, \ldots, C_l$ are disjoint connected critical submanifolds.

Definition 7. *The Morse-Bott polynomial of f is defined to be*

$$MB_t(f) = \sum_{j=1}^{l} P_t(C_j)t^{\lambda_j}$$

where λ_j is the Morse-Bott index of the critical submanifold C_j and $P_t(C_j)$ is the Poincaré polynomial of C_j.

Note: Clearly $MB_t(f)$ reduces to $M_t(f)$ when f is a Morse function. Also, if M or any of the critical submanifolds are not orientable, then $\mathbb{Z}_2$ coefficients are used to compute all the Betti numbers.

The following result is due to Bott. In fact, Bott proved a more general version of this result (without the assumption that the critical submanifolds are orientable) using homology with local coefficients in an orientation bundle [77, 7] [88, 8].

Theorem 8 (Morse-Bott Inequalities)**.** *Let $f : M \to \mathbb{R}$ be a Morse-Bott function on a finite dimensional oriented compact smooth manifold, and assume that all the critical submanifolds of f are orientable. Then there exists a polynomial $R(t)$ with non-negative integer coefficients such that*

$$MB_t(f) = P_t(M) + (1 + t)R(t).$$

If either M or some of the critical submanifolds are not orientable, then this equation holds when $\mathbb{Z}_2$ coefficients are used to define the Betti numbers.

5. A PERTURBATION TECHNIQUE AND THE PROOF OF THE MAIN THEOREM

To prove Theorem 8 we will use the following perturbation technique based on [22, 2], the Morse-Bott Lemma, and a folk theorem proved in [11, 1]. This construction produces an explicit Morse-Smale function $h : M \to \mathbb{R}$ arbitrarily close to a given Morse-Bott function $f : M \to \mathbb{R}$ such that $h = f$ outside of a neighborhood of the critical set $\mathrm{Cr}(f)$.

Let T_j be a small tubular neighborhood around each connected component $C_j \subseteq \mathrm{Cr}(f)$ for all $j = 1, \ldots, l$ with local coordinates (u, v, w) consistent with those from the Morse-Bott Lemma (Lemma 5). By "small" we mean that each T_j is contained in the union of the domains of the charts from the Morse-Bott Lemma, for $i \neq j$ we have $T_i \cap T_j = \emptyset$ and f decreases by at least three times $\max\{\mathrm{var}(f, T_j) |\ j = 1, \ldots, l\}$ along any gradient flow line from T_i to T_j where $\mathrm{var}(f, T_j) = \sup\{f(x) |\ x \in T_j\} - \inf\{f(x) |\ x \in T_j\}$, and if $f(C_i) \neq f(C_j)$, then

$$\mathrm{var}(f, T_i) + \mathrm{var}(f, T_j) < \frac{1}{3}\left| f(C_i) - f(C_j) \right|.$$

Pick a Riemannian metric on M such that the charts from the Morse-Bott Lemma are isometries with respect to the standard Euclidean metric on $\mathbb{R}^m$, and then pick positive Morse functions $f_j : C_j \to \mathbb{R}$ that are Morse-Smale with respect to the restriction of the Riemannian metric to C_j for all $j = 1, \ldots, l$. The Morse-Smale functions $f_j : C_j \to \mathbb{R}$ exist by the Kupka-Smale Theorem (see for instance Theorem 6.6. of [33, 3]).

For every $j = 1, \ldots, l$ extend f_j to a function on T_j by making f_j constant in the direction normal to C_j, i.e. f_j is constant in the v and w coordinates coming from the Morse-Bott Lemma. Let $\tilde{T}_j \subset T_j$ be a smaller tubular neighborhood of C_j with the same coordinates as T_j, and let ρ_j be a smooth nonincreasing bump function which is constant in the u coordinates, equal to 1 on $\tilde{T}_j$, and equal to 0 outside of T_j. Choose $\varepsilon > 0$ small enough so that

$$\sup_{T_j - \tilde{T}_j} \varepsilon \|\nabla \rho_j f_j\| < \inf_{T_j - \tilde{T}_j} \|\nabla f\|$$

for all $j = 1, \ldots l$, and define

$$h = f + \varepsilon \left(\sum_{j=1}^{l} \rho_j f_j \right).$$

The function $h : M \to \mathbb{R}$ is a Morse function close to f, and the critical points of h are exactly the critical points of the f_j for $j = 1, \ldots, l$. Moreover, if $p \in C_j$ is a critical point of $f_j : C_j \to \mathbb{R}$ of index λ_p^j, then p is a critical point of h of index $\lambda_p^h = \lambda_j + \lambda_p^j$.

We now apply a well-known folk theorem (whose proof can be found in Section 2.12 of [11, 1]) which says that we can perturb the Riemannian metric on M outside of the union of the tubular neighborhoods T_j for $j = 1, \ldots, l$ so that h satisfies the Morse-Smale transversality condition with respect to the perturbed metric. In summary, we have achieved the following conditions.

1. The gradient $\nabla f = \nabla h$ outside of the union of the tubular neighborhoods T_j for $j = 1, \ldots, l$. Moreover, the tubular neighborhood T_j are chosen small enough so that the following condition holds: if $f(C_i) \leq f(C_j)$ for some $i \neq j$, then there are no gradient flow lines of h from C_i to C_j.
2. The charts from the Morse-Bott Lemma are isometries with respect to the metric on M and the standard Euclidean metric on $\mathbb{R}^m$.
3. In the local coordinates (u, v, w) of a tubular neighborhood T_j we have $f = f(C) - |v|^2 + |w|^2$, ρ_j depends only on the v and w coordinates, and f_j depends only on the u coordinates. In particular, $\nabla f \perp \nabla f_j$ on T_j by the previous condition.
4. The function $h = f + \varepsilon f_j$ on the tubular neighborhood $\tilde{T}_j$.
5. The gradient ∇f dominates $\varepsilon \nabla \rho_j f_j$ on $T_j - \tilde{T}_j$.
6. The functions $h : M \to \mathbb{R}$ and $f_j : C_j \to \mathbb{R}$ satisfy the Morse-Smale transversality condition for all $j = 1, \ldots, l$.
7. For every $n = 0, \ldots, m$ we have the following description of the n^{th} Morse-Smale-Witten chain group of h in terms of the Morse-Smale-Witten chain

groups of the f_j for $j = 1, \dots, l$.

$$C_n(h) = \bigoplus_{\lambda_j + k = n} C_k(f_j)$$

Now, let ∂_*^h denote the Morse-Smale-Witten boundary operator of h, and let $\partial_*^{f_j}$ denote the Morse-Smale-Witten boundary operator of f_j for $j = 1, \dots, l$.

Lemma 9. *If $p, q \in C_j$ are critical points of $f_j : C_j \to \mathbb{R}$ of relative index one, then the coefficients $n(q,p)$ used to define the Morse-Smale-Witten boundary operator are the same for ∂_*^h and $\partial_*^{f_j}$ (assuming in the case where C_j is orientable that the appropriate orientation of C_j has been chosen).*

Proof: We have $h = f + \varepsilon f_j$ on $\tilde{T}_j$ by condition (4). This implies that $\nabla h = \nabla f + \varepsilon \nabla f_j$ on $\tilde{T}_j$, and $\nabla h = \varepsilon \nabla f_j$ on the critical submanifold C_j. Moreover, by conditions (3) and (5) a gradient flow line of h cannot begin and end in C_j unless the entire flow line is contained in C_j. Thus, if p and q are both in C_j the flows connecting them along the gradient flow lines of h and of f_j are the same, and the numbers $n(q,p)$ in the complex of h and of f_j are the same as long as the orientations are chosen appropriately.

□

We now order the connected critical submanifolds $C_1, \dots, C_l$ by height, i.e. such that $f(C_i) \leq f(C_j)$ whenever $i \leq j$. For $\alpha \in C_n(h)$ and C_j a connected critical submanifold we denote by α_j the chain obtained from α by retaining only those critical points belonging to C_j. By condition (7), any $\alpha \neq 0$ can be written uniquely as:

$$\alpha = \alpha_{j_1} + \cdots + \alpha_{j_r}$$

where $j_1 < \cdots < j_r$ and $\alpha_{j_i} \in C_{k_i}(f_{j_i}) - \{0\}$ where $\lambda_{j_i} + k_i = n$. We will refer to α_{j_r} as the "top part" of the chain α.

Corollary 10. *If α is a non-zero element in ker ∂_*^h, then the top part of α is a non-zero element in the kernel of $\partial_*^{f_j}$ for some j, i.e. using the above notation for $\alpha \in$ ker ∂_n^h we have $\alpha_{j_r} \in$ ker $\partial_{k_r}^{f_{j_r}}$ where $\lambda_{j_r} + k_r = n$.*

Proof: Let $\alpha = \sum_i n_i q_i \in \ker \partial_n^h$, i.e.

$$0 = \partial_n^h(\alpha) = \sum_i n_i \partial_n^h(q_i) = \sum_i n_i \sum_{p \in Cr_{n-1}(h)} n(q_i, p) p = \sum_{p \in Cr_{n-1}(h)} \left(\sum_i n_i n(q_i, p) \right) p.$$

Then $\sum_i n_i n(q_i, p) = 0$ for all $p \in Cr_{n-1}(h)$, and in particular, for $p \in Cr_{n-1}(h) \cap C_{j_r}$. For any $p \in Cr_{n-1}(h) \cap C_{j_r}$ we have $f(q_i) \leq f(p)$ for all q_i in the sum of α such that

$q_i \notin C_{j_r}$, and thus there are no gradient flow lines of h from q_i to p by condition (1). Therefore, $n(q_i, p) = 0$ if $p \in Cr_{n-1}(h) \cap C_{j_r}$ and $q_i \notin C_{j_r}$, i.e.

$$0 = \sum_i n_i n(q_i, p) = \sum_{q_i \in C_{j_r}} n_i n(q_i, p)$$

when $p \in Cr_{n-1}(h) \cap C_{j_r}$. Lemma 9 then implies that $\partial_{k_r}^{f_{j_r}}(\alpha_{j_r}) =$

$$\sum_{q_i \in C_{j_r}} n_i \partial_{k_r}^{f_{j_r}}(q_i) = \sum_{q_i \in C_{j_r}} n_i \sum_{p \in Cr_{k_r-1}(f_{j_r})} n(q_i, p)p = \sum_{p \in Cr_{k_r-1}(f_{j_r})} \left(\sum_{q_i \in C_{j_r}} n_i n(q_i, p) \right) p = 0,$$

where $\lambda_{j_r} + k_r = n$ and $Cr_{k_r-1}(f_{j_r}) = Cr_{n-1}(h) \cap C_{j_r}$.

□

Proof of Theorem 8: Let f, h, f_j for $j = 1, \ldots, l$, and the Riemannian metric on M be as above. Let $M_t(f_j)$ denote the Morse polynomial of $f_j : C_j \to \mathbb{R}$, and let $c_j = \dim C_j$ for all $j = 1, \ldots, l$. Note that the relation $\lambda_p^h = \lambda_j + \lambda_p^j$ implies that

$$M_t(h) = \sum_{j=1}^{l} M_t(f_j) t^{\lambda_j}.$$

The polynomial Morse inequalities (Theorem 3) say that

$$M_t(h) = P_t(M) + (1+t)R_h(t)$$

and

$$M_t(f_j) = P_t(C_j) + (1+t)R_j(t)$$

where $R_h(t)$ and $R_j(t)$ are polynomials with non-negative integer coefficients for all $j = 1, \ldots, l$.

$$\begin{aligned}
MB_t(f) &= \sum_{j=1}^{l} P_t(C_j) t^{\lambda_j} \\
&= \sum_{j=1}^{l} \left(M_t(f_j) - (1+t)R_j(t) \right) t^{\lambda_j} \\
&= \sum_{j=1}^{l} M_t(f_j) t^{\lambda_j} - (1+t) \sum_{j=1}^{l} R_j(t) t^{\lambda_j} \\
&= M_t(h) - (1+t) \sum_{j=1}^{l} R_j(t) t^{\lambda_j} \\
&= P_t(M) + (1+t)R_h(t) - (1+t) \sum_{j=1}^{l} R_j(t) t^{\lambda_j}
\end{aligned}$$

In the proof of the polynomial Morse inequalities we saw that

$$R_j(t) = \sum_{k=1}^{c_j} (\nu_k^j - z_k^j) t^{k-1}$$

where ν_k^j is the rank of the group $C_k(f_j)$ and z_k^j is the rank of the kernel of the boundary operator $\partial_k^{f_j} : C_k(f_j) \to C_{k-1}(f_j)$ in the Morse-Smale-Witten chain complex of $f_j : C_j \to \mathbb{R}$. Hence,

$$MB_t(f) = P_t(M) + (1+t)\sum_{n=1}^{m} (\nu_n^h - z_n^h) t^{n-1} - (1+t)\sum_{j=1}^{l}\sum_{k=1}^{c_j} (\nu_k^j - z_k^j) t^{\lambda_j + k - 1}$$

where ν_n^h denotes the rank of the chain group $C_n(h)$ and z_n^h denotes the rank of the kernel of the Morse-Smale-Witten boundary operator $\partial_n^h : C_n(h) \to C_{n-1}(h)$. Since the critical points of h coincide with the critical points of the functions f_j for $j = 1, \ldots l$, and a critical point $p \in C_j$ of f_j of index λ_p^j is a critical point of h of index $\lambda_j + \lambda_p^j$ we have

$$\sum_{n=1}^{m} (\nu_n^h - z_n^h) t^{n-1} - \sum_{j=1}^{l}\sum_{k=1}^{c_j} (\nu_k^j - z_k^j) t^{\lambda_j + k - 1} = \sum_{j=1}^{l}\sum_{k=1}^{c_j} z_k^j t^{\lambda_j + k - 1} - \sum_{n=1}^{m} z_n^h t^{n-1}.$$

Therefore,

$$MB_f(t) = P_t(M) + (1+t)R(t)$$

where

$$R(t) = \sum_{j=1}^{l}\sum_{k=1}^{c_j} z_k^j t^{\lambda_j + k - 1} - \sum_{n=1}^{m} z_n^h t^{n-1} = \sum_{n=1}^{m} \left(\sum_{\lambda_j + k = n} z_k^j - z_n^h \right) t^{n-1}.$$

To see that $\sum_{\lambda_j + k = n} z_k^j \geq z_n^h$ for all $n = 1, \ldots, m$ we apply Corollary 10 as follows. If $z_n^h > 0$, then we can choose a nonzero element $\beta_1 \in \ker\, \partial_n^h$. By Corollary 10, the "top part" of β_1 is a nonzero element in $\ker\, \partial_k^{f_j}$ for some k and j such that $\lambda_j + k = n$. If $z_n^h = 1$, then we are done. If $z_n^h > 1$, then we can find an element $\beta_2 \in \ker\, \partial_n^h$ that is not in the group generated by β_1, and by adding a multiple of β_1 to β_2 if necessary, we can choose β_2 such that the top part of β_2 is a nonzero element that is not in the subgroup generated by the top part of β_1. Continuing in this fashion we can find generators for $\ker\, \partial_n^h$ whose top parts are independent elements in

$$\bigoplus_{\lambda_j + k = n} \ker\, \partial_k^{f_j},$$

i.e.

$$\sum_{\lambda_j + k = n} \left(\text{rank } \ker\, \partial_k^{f_j} \right) \geq \text{rank } \ker\, \partial_n^h.$$

□

Acknowledgments: We would like to thank William Minicozzi for several insightful conversations concerning Riemannian metrics and gradient flows.

References

[1]]cite.AbbLec1Alberto Abbondandolo and Pietro Majer, *Lectures on the Morse complex for infinite-dimensional manifolds*, Morse theoretic methods in nonlinear analysis and in symplectic topology, 2006, pp. 1–74. MR2276948

[2]]cite.AusMor2D. M. Austin and P. J. Braam, *Morse-Bott theory and equivariant cohomology*, The Floer memorial volume, 1995, pp. 123–183. MR1362827 (96i:57037)

[3]]cite.BanLec3Augustin Banyaga and David Hurtubise, *Lectures on Morse homology*, Kluwer Texts in the Mathematical Sciences, vol. 29, Kluwer Academic Publishers Group, Dordrecht, 2004. MR2145196 (2006i:58016)

[4]]cite.BanApr4Augustin Banyaga and David E. Hurtubise, *A proof of the Morse-Bott lemma*, Expo. Math. **22** (2004), no. 4, 365–373. MR2075744 (2005b:57062)

[5]]cite.BanMor5——, *Morse-Bott homology* (2006), available at `arXiv:math.AT/0612316`.

[6]]cite.BisThe6Jean-Michel Bismut, *The Witten complex and the degenerate Morse inequalities*, J. Differential Geom. **23** (1986), no. 3, 207–240. MR852155 (87m:58169)

[7]]cite.BotNon7Raoul Bott, *Nondegenerate critical manifolds*, Ann. of Math. (2) **60** (1954), 248–261. MR0064399 (16,276f)

[8]]cite.BotLec8——, *Lectures on Morse theory, old and new*, Bull. Amer. Math. Soc. (N.S.) **7** (1982), no. 2, 331–358. 84m:58026a

[9]]cite.BotMor9——, *Morse theory indomitable*, Inst. Hautes Études Sci. Publ. Math. (1988), no. 68, 99–114 (1989). 90f:58027

[10]]cite.HelApr10B. Helffer and J. Sjöstrand, *A proof of the Bott inequalities*, Algebraic analysis, vol. i, 1988, pp. 171–183. MR992453 (90g:58018)

[11]]cite.HenLes11Guy Henniart, *Les inégalités de Morse (d'après E. Witten)*, Astérisque (1985), no. 121-122, 43–61. Séminaire Bourbaki, Vol. 1983/84. MR768953 (86f:58027)

[12]]cite.JiaMor12Mei-Yue Jiang, *Morse homology and degenerate Morse inequalities*, Topol. Methods Nonlinear Anal. **13** (1999), no. 1, 147–161. 2000f:57037

[13]]cite.SchMor13Matthias Schwarz, *Morse homology*, Birkhäuser Verlag, Basel, 1993. MR1239174 (95a:58022)

[14]]cite.WitSup14Edward Witten, *Supersymmetry and Morse theory*, J. Differential Geom. **17** (1982), no. 4, 661–692 (1983). 84b:58111

Department of Mathematics, Penn State University, University Park, PA 16802
E-mail address: `banyaga@math.psu.edu`

Department of Mathematics and Statistics, Penn State Altoona, Altoona, PA 16601-3760
E-mail address: `Hurtubise@psu.edu`